

Suggested French Bulldog Anesthesia Protocol

Lori Hunt, DVM

Requirement:

No food for 12 hours before surgery. This is crucial. If the dog eats the morning of surgery, cancel the surgery for that day. Vomiting and aspiration can have devastating effects in a Frenchie.

Recommendations:

- 1) Do a clotting time, full blood chemistry work-up and complete blood count before anesthesia.
- 2) Be sure an intravenous catheter will be placed prior to surgery.
- 3) Ideally, all dogs will be administered IV fluids during surgery.
- 4) Chest x-rays prior to surgery are always recommended by me for brachycephalic dogs, especially if the dog has had chronic breathing problems.

Optimum

Use propofol induction anesthesia, intubate (place a breathing tube in the trachea) and maintain on gas anesthesia (isoflurane or sevoflurane).

Satisfactory Options

Ketamine combined with diazepam (Valium)

Butorphanol (mild sedative for short procedures such as an x-ray) This is also called Torbugesic or Torbutrol

Dexdomitor (reversible anesthesia/sedative).

Use with Caution

Thiopental

Telazol

Do Not Give Frenchies

Acepromazine (sedative)

Pentobarbital (injectable anesthesia)

Xylazine (sedative)

Halothane (gas anesthesia)

Anesthetic Extras

(These would be in addition to the optimum anesthetic protocol listed above and are recommended for longer procedures)

Famotidine (Pepcid) or ranitidine (Zantac) injection (helps cut down on nausea and post-op vomiting, decreases risk of aspiration)

Dexamethasone can reduce post-op swelling and make recovery faster in cases where the dog's throat is irritated from the ET tube, when the dog's palate is very long, or following palate surgery. A single injection can be given in these cases.

Intubation vs. Masking/Coning Down

EVERY brachycephalic dog that goes under anesthesia should have an endotracheal tube (ET) placed in his or her trachea. Always! The airway must be protected at all times. The endotracheal tube should be left in until the dog is awake. Use intravenous propofol (or one of the other satisfactory drugs listed above) to induce anesthesia (which puts them under) and allows sufficient time to place the ET tube. From then on, anesthesia is maintained with sevoflurane or isoflurane.

Brachycephalic breeds, such as Frenchie, should never be masked down with anesthesia. Masking down is when a mask is placed on an awake or mildly sedated dog's muzzle. The mask is held in place by restraining the dog. The inhalant, which has a bad smell, is given at high levels. As the dog breaths more and more, he/she gets sleepy. The problem with masking down Frenchies is that they can become very anxious, fight the mask and not breath well. Most Frenchies have problems breathing in the first place, this just makes it worse, which results in lower oxygen levels. Ideally, injectable sedatives are used and an endotracheal tube placed which is then attached to an anesthetic machine. This gives them the optimum oxygen supply.

Rev 3/13